


# Various tools are available to "downscale" regional and local climate change information




### "Nested" Regional Climate Modeling

Motivation: The resolution of GCMs is still too coarse to capture regional and local climate processes

Technique: A "Regional Climate Model" (RCM) is "nested" within a GCM in order to locally increase the model resolution.


 Initial conditions (IC) and lateral boundary conditions (LBC) for the RCM are obtained from the GCM ("One-way Nesting") or analyses of observations.

Strategy: The GCM simulates the response of the general circulation to the large scale forcings, the RCM simulates the effect of sub-GCM-grid scale forcings and provides fine scale regional information




Technique borrowed from NWP


Number of papers in the ISI under "regional climate model"




### Some key projects and literature

- Review papers: Giorgi and Mearns (1991), McGregor (1997), Giorgi and Mearns (1999), Giorgi et al. (IPCC 2001), Leung et al. (2003), Mearns et al. (2003), Wang et al. (2004), Giorgi (2006), Laprise (2007)
- European projects: PRUDENCE, AMMA, ENSEMBLES, CECILIA, CLARIS, ACQWA
- Intercomparison projects: PIRCS, RMIP, NARCCAP, NEWBALTIC, ARCMIP, PLATIN, ARC, NAMAP, QUIRCS, Transferability
- Special issues: JGR 1999; JMSJ 2004; TAC 2006; CC 2007; MAP 2004, 2008; CCH 2006; MET.-ZEIT. 2008







#### **Dynamical Downscaling**


Generation of small scales by a high-resolution RCM driven by low-resolution GCM data (900 hPa specific humidity)

**Large scales Short scales** 

From Laprise et al. (2007)

## RCM simulation of precipitation at different resolutions over the Alps (1960-1990)


Mean annual precipitation (mm/day)




From Christensen et al. 2005


### The added value of RCMs


#### **Topographic forcing**






Courtesy of R. Jones









DJF precipitation 30-year nested RCM simulation, 1961-1990, 20 km grid spacing


#### **OBSERVATIONS**



Simulation of east Asia monsoon precipitation by GCMs has been traditionally very difficult


(Mean annual precipitation is shown; Gao et al. 2006)








### Mean annual precipitation (mm/day)





# Simulation of mean precipitation for an all-Africa domain


## ERA-Interim LBC (1989-2007)

Sylla et al. 2009


a. West Sahel Precip b. East Sahel Precip c. Guinea Coast Precip d. North-Equatorial Central-Africa Precip e. Horn Africa Precip f. South-Equatorial Central-Africa Precip JÁN FÉB MÁR APR MÁY JÚN JÚL AÚG SÉP OCT NÓV ĐỂC g. Central South-Africa Precip h. South Africa Precip FEB MAR APR MAY JUN JUL AUG SEP MÁR APR MÁY JÚN JÚL AUG SÉP

# Simulation of seasona precipitation for an all-Africa domain

## ERA-Interim LBC (1989-2007)




Sylla et al. 2009




# Simulation of interannual variability for an all-Africa domain

## ERA-Interim LBC (1989-2007)




Sylla et al. 2009

### Surface winds in a coupled regional RCM for the Mediterranean 30 km resolution, 1960-1990, ERA40 LBC (Artale et al. 2009)




## Do high resolution forcings affect the climate change signal? Topography

DJF Precipitation change (%)- dx=20 km A2 (2071-2100) – Control (1961-1990) (Gao et al. 2006)




# Do high resolution forcings affect the climate change signal? Coastlines


### Summer temperature change, 2080s, A2





(Courtesy of R. Jones)

## Do high resolution forcings affect the climate change signal? Aerosols











### Regional Climate Models: "State of the art"


- Many RCMs today available, some of them "portable" and used by wide communities (e.g. RegCM, PRECIS, RSM, WRF)
- Grid spacing of 10-30 km (sub-10 km for some models);
- Upgrade to non-hydrostatic, cloud-resolving frameworks in order to go to sub-10 km resolutions
- Decadal to centennial simulations the "accepted standard"
- Virtually all regions of the World have been simulated
- Encouraging results from some two-way nested experiments
- Several RCM coupling efforts under way, including atmosphere, ocean, aerosol, and biosphere components
- Wide range of applications
  - Process studies, paleoclimate, climate change, seasonal prediction, impacts, climate-aerosol interactions, air-sea interactions, landatmosphere feedbacks



# There are many sources of uncertainty to regional climate change projection




#### Regional Climate Change "Hyper-Matrix Framework" (HMF)




# CORDEX: COordinated Regional climate Downscaling Experiment

- Task Force on Regional Climate Downscaling (TFRCD)
  - Evaluate and possibly improve different regional downscaling techniques (Model Evaluation Framework, AMIP-like)
  - Design a common framework for the next generation RCD-based climate change projections for input to impact/adaptation work (Model Projection Framework, CMIP-like)
  - Facilitate the engagement of the end-user community and the scientific community from developing countries


### The CORDEX Framework



### **CORDEX time slices**



### CORDEX Phase I experiment design



Regional Analysis
Regional Databanks

1951-2100 1981-2010, 2041-2070, 2011-2040

 Regional climate modeling (or more generally downscaling) should not be viewed as alternative to global high resolution modeling, but as complimentary

- Regional climate modeling (or more generally downscaling) should not be viewed as alternative to global high resolution modeling, but as complimentary
- RCMs are downscaling tools, so are affected by the forcing fields. It should not be asked of them more than they can do.


- Regional climate modeling (or more generally downscaling) should not be viewed as alternative to global high resolution modeling, but as complimentary
- RCMs are downscaling tools, so are affected by the forcing fields. It should not be asked of them more than they can do.
- RCMs and other downscaling techniques can be valuable tools to study regional processes and interface with the end-user community

- Regional climate modeling (or more generally downscaling) should not be viewed as alternative to global high resolution modeling, but as complimentary
- RCMs are downscaling tools, so are affected by the forcing fields. It should not be asked of them more than they can do.
- RCMs and other downscaling techniques can be valuable tools to study regional processes and interface with the end-user community
- RCMs can enlarge the modeling community and as such have an important "educational" role


- Regional climate modeling (or more generally downscaling) should not be viewed as alternative to global high resolution modeling, but as complimentary
- RCMs are downscaling tools, so are affected by the forcing fields. It should not be asked of them more than they can do.
- RCMs and other downscaling techniques can be valuable tools to study regional processes and interface with the end-user community
- RCMs can enlarge the modeling community and as such have an important "educational" role
- CORDEX will provide an important benchmark framework for the downscaling community.



#### Climate simulation segment of the uncertainty cascade




#### Cascade of uncertainty in climate change projection




### Simulation of extreme precipitation


Summer 1993 Flood




Summer 1988 Drought



Summer 1993 Flood



Summer 1988 Drought



#### JJA Precipitation

